

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
4 January 2001 (04.01.2001)

PCT

(10) International Publication Number
WO 01/00659 A1

- (51) International Patent Classification⁷: C07K 5/06, (74) Agents: WOODROW, Hal, Brent et al.; Johnson & Johnson, One Johnson & Johnson Plaza, New Brunswick, NJ 08903 (US).
C07D 235/04
- (21) International Application Number: PCT/US00/17751
- (22) International Filing Date: 28 June 2000 (28.06.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/141,552 29 June 1999 (29.06.1999) US
09/599,826 22 June 2000 (22.06.2000) US
- (71) Applicants: ORTHO-MCNEIL PHARMACEUTICAL, INC. [US/US]; U.S. Route #202, P.O. Box 300, Raritan, NJ 08869-0602 (US). COR THERAPEUTICS, INC. [US/US]; 256 East Grand Avenue, South San Francisco, CA 94080 (US).
- (72) Inventors: ZHANG, Han-Cheng; 109 Bryan Way, Lansdale, PA 19446 (US). MARYANOFF, Bruce, E.; 4029 Devonshire Drive, Forest Grove, PA 18922 (US). MC-COMSEY, David, F.; 1125 Victoria Road, Warminster, PA 18974 (US). WHITE, Kimberly, B.; 120 Blue Ribbon Drive, North Wales, PA 19454 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— With international search report.
— Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: BENZIMIDAZOLONE PEPTIDOMIMETICS AS THROMBIN RECEPTOR ANTAGONISTS

(57) Abstract: The invention is directed to novel benzimidazolone peptidomimetic compounds which are useful as thrombin receptor antagonists for the treatment of diseases associated with thrombosis, restenosis, hypertension, heart failure, arrhythmia, inflammation, angina, stroke, atherosclerosis, ischemic conditions, Angiogenesis related disorders, cancer, and neurodegenerative disorders. Pharmaceutical compositions comprising the substituted benzimidazolone peptidomimetics of the present invention and methods of treating conditions mediated by the thrombin receptor are also disclosed.

WO 01/00659 A1

BENZIMIDAZOLONE PEPTIDOMIMETICS AS THROMBIN RECEPTOR ANTAGONISTS

5 FIELD OF THE INVENTION

This patent application claims priority from provisional patent application Serial Number 60/141,552, which was filed June 29, 1999. This invention relates to certain novel thrombin receptor antagonists, their synthesis and their
10 use for the treatment of diseases associated with thrombosis, restenosis, hypertension, heart failure, arrhythmia, inflammation, angina, stroke, atherosclerosis, ischemic conditions, Angiogenesis related disorders, cancer, and neurodegenerative disorders.

15 BACKGROUND OF THE INVENTION

Thrombin is an important serine protease in hemostasis and thrombosis. One of the key actions of thrombin is cellular modulation via receptor activation. A functional human thrombin receptor (PAR-1), cloned by Coughlin
20 in 1991 (T.-K. Vu, *Cell* **1991**, 64, 1057), was found to be a member of the G-protein coupled receptor (GPCR) superfamily. The receptor activation putatively occurs by N-terminal recognition and proteolytic cleavage at the Arg-41/Ser-42 peptide bond to reveal a truncated N-terminus. This new receptor sequence, which has an SFLLRN (Ser-Phe-Leu-Leu-Arg-Asn) N-terminus
25 acting as a tethered ligand to recognize a site on the receptor, can trigger activation and signal transduction leading to platelet aggregation. Since 1991, three other protease-activated receptors with extensive homology to the thrombin receptor, "PAR-2" (S. Nystedt, *Proc. Natl. Acad. Sci USA* **1994**, 91, 9208), "PAR-3" (H. Ishihara, *Nature* **1997**, 386, 502), and "PAR-4" (W.-F. Xu,
30 *Proc. Natl. Acad. Sci USA* **1998**, 95, 6642), have been cloned. Thrombin receptor (PAR-1) specific antibody-induced blockade of the platelet thrombin receptor has shown efficacy against arterial thrombosis in vivo (J. J. Cook *Circulation* **1995**, 91, 2961). Hence, antagonists of the thrombin receptor (PAR-1) are useful to block these protease-activated receptors and, as such,
35 may be used to treat platelet mediated thrombotic disorders such as


myocardial infarction, stroke, restenosis, angina, atherosclerosis, and ischemic conditions.

The thrombin receptor (PAR-1) has also been identified on other cell types: endothelial, fibroblast, renal, osteosarcoma, smooth muscle, myocytes, tumor, and neuronal/glia. Thrombin activation of endothelial cells upregulates P-selectin to induce polymorphonuclear leukocyte adhesion - an inflammatory response of the vessel wall (Y. Sugama, *J. Cell Biol.* 1992, 119, 935). In fibroblasts, thrombin receptor (PAR-1) activation induces proliferation and transmission of mitogenic signals (D. T. Hung, *J. Cell Biol.* 1992, 116, 827). Thrombin has been implicated in osteoblast proliferation through its activation of osteoblast cells (D. N. Tatakis, *Biochem. Biophys. Res. Commun.* 1991, 174, 181). Thrombin has been implicated in the regulation and retraction of neurons (K. Jalink, *J. Cell. Biol.* 1992, 118, 411). Therefore, in this context, the antagonist compounds of this invention may also be useful against inflammation, osteoporosis, Angiogenesis related disorders, cancer, neurodegenerative disorders, hypertension, heart failure, arrhythmia, glomerulonephritis.

The compounds of the present invention are a structurally novel class of benzimidazolone peptidomimetics represented by the general formula (I) below.

SUMMARY OF THE INVENTION

The present invention is directed to structurally novel compounds represented by the following general formula (I):



(I)

wherein

- 5 A_1 and A_2 are each independently a D- or L-amino acid selected from the group consisting of alanine, β -alanine, arginine, homoarginine, cyclohexylalanine, citrulline, cysteine (optionally substituted with C_1 - C_4 alkyl, aryl, or arC_1 - C_4 alkyl), 2,4-diaminobutyric acid (optionally substituted with acyl, C_1 - C_4 alkyl, aroyl, amidino, or $MeC(NH)-$), 2,3 diaminopropionic acid (optionally substituted with acyl, C_1 - C_4 alkyl, aroyl, amidino, or $MeC(NH)-$), glutamine, 10 glycine, indanylglycine, lysine (optionally substituted with acyl, C_1 - C_4 alkyl, aroyl, $MeC(NH)-$), valine, methionine, proline, serine (optionally substituted with C_1 - C_4 alkyl, aryl, or arC_1 - C_4 alkyl), homoserine (optionally substituted with C_1 - C_4 alkyl, aryl, or arC_1 - C_4 alkyl), tetrahydroisoquinoline-3-COOH, threonine 15 (optionally substituted with C_1 - C_4 alkyl, aryl, or arC_1 - C_4 alkyl), ornithine (optionally substituted with acyl, C_1 - C_4 alkyl, aroyl, $MeC(NH)-$), and an unsubstituted or substituted aromatic amino acid selected from the group consisting of phenylalanine, heteroarylalanine, naphthylalanine, homophenylalanine, histidine, tryptophan, tyrosine, arylglycine, 20 heteroarylglycine, aryl- β -alanine, and heteroaryl- β -alanine wherein the substituents on the aromatic amino acid are independently selected from one or more of halogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, hydroxy, C_1 - C_4 alkoxycarbonyl, amino, amidino, guanidino, fluorinated C_1 - C_4 alkyl, fluorinated C_1 - C_4 alkoxy, C_1 - C_4 alkylsulfonyl, C_1 - C_4 alkylcarbonyl, cyano, aryl, heteroaryl, arC_1 - C_4 alkyl, C_2 - C_4 alkenyl, alkynyl, or nitro; 25

Preferably, A_1 and A_2 are each independently an L-amino acid selected from the group consisting of alanine, β -alanine, arginine, homoarginine, cyclohexylalanine, citrulline, cysteine (optionally substituted with C_1 - C_4 alkyl, aryl, or arC_1 - C_4 alkyl), 2,4-diaminobutyric acid (optionally substituted with acyl, C_1 - C_4 alkyl, aroyl, amidino, or $MeC(NH)-$), 2,3 diaminopropionic acid (optionally substituted with acyl, C_1 - C_4 alkyl, aroyl, amidino, or $MeC(NH)-$), glutamine, glycine, indanylglycine, lysine (optionally substituted with acyl, C_1 - C_4 alkyl, 30

aroyl, MeC(NH)-), valine, methionine, proline, serine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), homoserine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), tetrahydroisoquinoline-3-COOH, threonine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), ornithine
 5 (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), and an unsubstituted or substituted aromatic amino acid selected from the group consisting of phenylalanine, heteroarylalanine, naphthylalanine, homophenylalanine, histidine, tryptophan, tyrosine, arylglycine, heteroarylglycine, aryl-β-alanine, and heteroaryl-β-alanine wherein the
 10 substituents on the aromatic amino acid are independently selected from one or more of halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, C₁-C₄ alkoxycarbonyl, amino-, amidino, guanidino, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl, heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or nitro;

15

R₁ is selected from amino, C₁-C₈ alkylamino, C₁-C₈ dialkylamino, arylamino, arC₁-C₈ alkylamino, C₃-C₈ cycloalkylamino, heteroalkylC₁-C₈ alkylamino, heteroalkylC₁-C₈ alkyl-N-methylamino, C₁-C₈ dialkylaminoC₁-C₈ alkylamino, -N(C₁-C₈alkyl)-C₁-C₈ alkyl-N(C₁-C₈alkyl)₂, N(C₁-C₈ alkyl)(C₁-C₈
 20 alkenyl), -N(C₁-C₈alkyl)(C₃-C₈cycloalkyl), heteroalkyl or substituted heteroalkyl wherein the substituent on the heteroalkyl is selected from oxo, amino, C₁-C₈ alkoxyC₁-C₈ alkyl, C₁-C₈ alkylamino or C₁-C₈ dialkylamino;

Preferably, R₁ is selected from amino, C₁-C₆ alkylamino, C₁-C₆ dialkylamino, arylamino, arC₁-C₆ alkylamino, heteroalkylC₁-C₆ alkylamino, -N(C₁-C₆alkyl)-C₁-C₆alkyl-N(C₁-C₆alkyl)₂, heteroalkyl or substituted heteroalkyl
 25 wherein the substituent on the heteroalkyl is selected from oxo, amino, C₁-C₆alkoxyC₁-C₆ alkyl, C₁-C₆ alkylamino or C₁-C₆ dialkylamino;

30 R₂ and R₃ are each independently selected from hydrogen, C₁-C₈ alkyl, C₃-C₈ cycloalkyl, C₃-C₈ cycloalkylC₁-C₈ alkyl, aryl, heteroalkyl, substituted heteroalkyl (wherein the substituent on the heteroalkyl is one or more substituents independently selected from C₁-C₈ alkoxycarbonyl, C₁-C₈ alkyl, or C₁-C₄ alkylcarbonyl), heteroalkylC₁-C₈ alkyl, indanyl, acetamidinoC₁-C₈ alkyl,
 35 aminoC₁-C₈ alkyl, C₁-C₈ alkylaminoC₁-C₈ alkyl, C₁-C₈ dialkylaminoC₁-C₈ alkyl,

unsubstituted or substituted heteroarylC₁-C₈ alkyl or unsubstituted or substituted arC₁-C₈ alkyl, wherein the substituent on the aralkyl or heteroarylalkyl group is one or more substituents independently selected from halogen, nitro, amino, C₁-C₈ alkyl, C₁-C₈ alkoxy, hydroxy, cyano, C₁-C₄ alkylcarbonyl, C₁-C₈ alkoxycarbonyl, hydroxyC₁-C₈ alkyl or aminosulfonyl; or

R₂ and R₃ together with the nitrogen to which they are attached, alternatively form an unsubstituted or substituted heteroalkyl group selected from piperidiny, piperaziny, morpholinyl or pyrrolidinyl, wherein the substituent is one or more substituents independently selected from C₁-C₈ alkyl C₁-C₈ alkoxycarbonyl or C₁-C₄ alkylcarbonyl;

Preferably, R₂ is selected from hydrogen or C₁-C₆ alkyl;

R₃ is selected from C₁-C₈ alkyl, C₃-C₆ cycloalkyl, C₃-C₆ cycloalkylC₁-C₆ alkyl, aryl, heteroarylC₁-C₆ alkyl, substituted heteroarylC₁-C₆ alkyl wherein the substituent is C₁-C₄ alkyl, heteroalkyl, heteroalkylC₁-C₆ alkyl, indanyl, acetamidinoC₁-C₈ alkyl, aminoC₁-C₆ alkyl, C₁-C₈ alkylaminoC₁-C₆ alkyl, C₁-C₆ dialkylaminoC₁-C₆ alkyl, arC₁-C₈ alkyl, substituted arC₁-C₈ alkyl wherein the substituent on the aralkyl group is one to five substituents independently selected from halogen, nitro, amino, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ alkoxycarbonyl, hydroxyalkyl or aminosulfonyl; or

R₂ and R₃, together with the nitrogen to which they are attached, alternatively form an unsubstituted or substituted heteroalkyl group selected from piperidiny, piperaziny or pyrrolidinyl, wherein the substituent is independently one or two substituents selected from C₁-C₆ alkyl;

R₄ is selected from unsubstituted or substituted aryl, arC₁-C₈ alkyl, C₃-C₈ cycloalkyl, or heteroaryl, where the substituents on the aryl, arC₁-C₈ alkyl, cycloalkyl or heteroaryl group are independently selected from one or more of halogen, nitro, amino, cyano, hydroxyalkyl, C₁-C₈ alkyl, C₁-C₈ alkoxy, hydroxy, C₁-C₄ alkylcarbonyl, C₁-C₈ alkoxycarbonyl, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy or C₁-C₄ alkylsulfonyl;

Preferably, R₄ is selected from unsubstituted or substituted aryl, arC₁-C₆ alkyl, C₃-C₆ cycloalkyl or heteroaryl, where the substituents on the aryl, aralkyl, cycloalkyl or heteroaryl group are independently selected from one to three

substituents selected from halogen, cyano, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ alkoxycarbonyl, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy or C₁-C₄ alkylsulfonyl;

5 R₅ is selected from hydrogen or C₁-C₈ alkyl; preferably, R₅ is hydrogen;

X is oxygen or sulfur; preferably, X is oxygen;

m is an integer selected from 0, 1, 2 or 3;

10 n is an integer selected from 2 or 3; and

p is an integer selected from 0 or 1; preferably, p is 1;

and pharmaceutically acceptable salts thereof.

15 In a preferred embodiment of the present invention:

A₁ is an L-amino acid selected from the group consisting of alanine, arginine, cyclohexylalanine, glycine, proline, tetrahydroisoquinoline-3-COOH, and an unsubstituted or substituted aromatic amino acid selected from the
20 group consisting of phenylalanine, naphthylalanine, homophenylalanine, and O-methyl tyrosine, wherein the substituents on the aromatic amino acid are independently selected from one to five of (preferably, one to three of) halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, C₁-C₄ alkoxycarbonyl, amino, amidino, guanidino, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄
25 alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl, heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or nitro;

A₂ is an L-amino acid selected from the group consisting of alanine, β-alanine, arginine, citrulline, cysteine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), 2,4-diaminobutyric acid (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, amidino, or MeC(NH)-), 2,3- diaminopropionic acid (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, amidino, or MeC(NH)-), glutamine, glycine, lysine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), valine, methionine, serine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), homoserine (optionally substituted with C₁-C₄ alkyl, aryl,
35

or arC₁-C₄ alkyl), threonine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), ornithine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), and an unsubstituted or substituted aromatic amino acid selected from the group consisting of phenylalanine, heteroarylalanine, and histidine,
5 wherein the substituents on the aromatic amino acid are independently selected from one to five of (preferably, one to three of) halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, C₁-C₄ alkoxycarbonyl, amino, amidino, guanidino, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl, heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or
10 nitro;

R₂ is selected from hydrogen or C₁-C₄ alkyl;

15 m is 1 and n is 2;

and all other variables are as defined previously;
and pharmaceutically acceptable salts thereof.


20 In a class of the invention:

A₁ is an L-amino acid selected from the group consisting of alanine, arginine, cyclohexylalanine, glycine, proline, and an unsubstituted or substituted aromatic amino acid selected from the group consisting of
25 phenylalanine, naphthylalanine, homophenylalanine, and O-methyl tyrosine, wherein the substituents on the aromatic amino acid are independently one to two substituents selected from halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, C₁-C₄ alkoxycarbonyl, amino, amidino, guanidino, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl,
30 heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or nitro;

A₂ is an L-amino acid selected from the group consisting of alanine, β-alanine, arginine, citrulline, cysteine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), 2,4-diaminobutyric acid (optionally substituted with acyl,

C₁-C₄ alkyl, aroyl, amidino, or MeC(NH)-, 2,3- diaminopropionic acid (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, amidino, or MeC(NH)-), glutamine, glycine, lysine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), valine, methionine, serine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), homoserine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), threonine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), ornithine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), and an unsubstituted or substituted aromatic amino acid selected from the group consisting of phenylalanine, heteroarylalanine, and histidine, wherein the substituents on the aromatic amino acid are independently one to two substituents selected from halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, C₁-C₄ alkoxycarbonyl, amino, amidino, guanidino, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl, heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or nitro;

R₁ is selected from dimethylamino, diethylamino, di-(*n*-propyl)amino,



Preferably, R₁ is: ;

R₂ is selected from hydrogen, methyl or ethyl;

R₃ is selected from 2- indanyl, phenyl, cyclohexylmethyl, cyclopentyl, pyridylmethyl, furanylmethyl, 2-(4-methyl-furanyl)methyl, thienylmethyl, diphenylmethyl, 4-imidazolylethyl, 2-(4-N-methyl)imidazolylethyl, *n*-octyl, phenyl-*n*-propyl, aminoethyl, aminopropyl, amino-*n*-pentyl, dimethylaminoethyl, 4-aminophenylsulfonylaminomethyl, acetamidineylethyl, 2-N-pyrrolidinylethyl, N-ethoxycarbonylpiperidinyl, unsubstituted or substituted phenylethyl or

unsubstituted or substituted benzyl wherein the substituents on the phenylethyl or benzyl are independently one or two substituents selected from methyl, fluorine, chlorine, nitro, methoxy, methoxycarbonyl or hydroxymethyl; or

- 5 R_2 and R_3 , together with the nitrogen to which they are attached, alternatively form a heteroalkyl group selected from piperidinyl, or 4-(N-methyl)piperazinyl;

- 10 R_4 is selected from cyclohexyl, 2-naphthyl, phenylethyl, 4-fluorophenylethyl, or unsubstituted or substituted phenyl, where the substituents on the phenyl are independently selected from one to two substituents selected from fluorine, chlorine, iodine, methyl, cyano, or trifluoromethyl;

- 15 Preferably, R_4 is 2,6-dichlorophenyl or 2-methylphenyl;

and all other variables are as defined previously;
and pharmaceutically acceptable salts thereof.

- 20 In a subclass of the invention,

A_1 is selected from 3,4-Difluorophenylalanine or 4-Chlorophenylalanine;

A_2 is selected from 2,4-Diaminobutyric acid or 4-Pyridylalanine;

R_2 is hydrogen;

- 25 R_3 is selected from benzyl or 2-aminoethyl;

and all other variables are as defined previously;
and pharmaceutically acceptable salts thereof.

- 30 Illustrative of the invention is a pharmaceutical composition comprising a pharmaceutically acceptable carrier and any of the compounds described above. Illustrating the invention is a pharmaceutical composition made by mixing any of the compounds described above and a pharmaceutically acceptable carrier. An illustration of the invention is a process for making a


pharmaceutical composition comprising mixing any of the compounds described above and a pharmaceutically acceptable carrier.

An example of the invention is a method of treating a disorder
5 (preferably, a platelet-mediated thrombotic disorder) selected from arterial and/or venous thrombosis, acute myocardial infarction, reocclusion following thrombolytic therapy and/or angioplasty, inflammation, unstable angina, stroke, restenosis, atherosclerosis, ischemic conditions, hypertension, heart failure, arrhythmia, glomerulonephritis, osteoporosis, Angiogenesis related disorders,
10 cancer, neurodegenerative disorders and a variety of vaso-occlusive disorders in a subject in need thereof comprising administering to the subject a therapeutically effective amount of any of the compounds or pharmaceutical compositions described above. In a preferred embodiment, the therapeutically effective amount of the compound is from about 0.1 mg/kg/day to about 300
15 mg/kg/day.


Also included in the invention is the use of any of the compounds described above for the preparation of a medicament for a disorder (preferably, a platelet-mediated thrombotic disorder) selected from arterial and/or venous
20 thrombosis, acute myocardial infarction, reocclusion following thrombolytic therapy and/or angioplasty, inflammation, unstable angina, stroke, restenosis, atherosclerosis, ischemic conditions, hypertension, heart failure, arrhythmia, glomerulonephritis, osteoporosis, Angiogenesis related disorders, cancer, neurodegenerative disorders or a variety of vaso-occlusive disorders in a
25 subject in need thereof.

DETAILED DESCRIPTION OF THE INVENTION

More particularly, the present invention is directed to compounds of the
30 following formula (I):



wherein A_1 , A_2 , R_1 , R_2 , R_3 , R_4 , R_5 , X , m , n and p are as previously defined. In a particularly preferred embodiment, the compounds have the formula




The compounds of the present invention are thrombin receptor antagonists and as such are useful in treating thrombosis, restenosis, hypertension, heart failure, arrhythmia, myocardial infarction, glomerulonephritis, reocclusion following thrombolytic therapy, reocclusion following angioplasty, inflammation, angina, stroke, atherosclerosis, ischemic conditions, a vaso-occlusive disorder, neurodegenerative disorders, Angiogenesis related disorders and cancer. These compounds are also useful as antithrombotics in conjunction with fibrinolytic therapy (e.g., t-PA or streptokinase).

In the compounds of formula (I), the amino acid residues comprising the A_1 and A_2 substituents are attached to the adjacent moiety according to standard nomenclature so that the amino-terminus (N-terminus) of the amino acid is drawn on the left and the carboxy-terminus of the amino acid is drawn on the right. So, for example, in Compound 1 in Table 1, where A_1 is 3,4-difluorophenylalanine and A_2 is Dbu (2,4-Diaminobutyric acid), the

N-terminus of the 3,4-difluorophenylalanine (A_1) is attached to the carbonyl group and the carboxy-terminus of the 3,4-difluorophenylalanine (A_1) is attached to the N-terminus of the A_2 substituent (Dbu), similarly, the N-terminus of the Dbu (A_2) is attached to the carboxy-terminus of the A_1 substituent and
 5 the carboxy-terminus of the Dbu (A_2) is attached to the $N-R_2R_3$ group.

When a particular group is "substituted" (e.g., Phe, aryl, heteroalkyl, heteroaryl), that group may have one or more substituents, preferably from one to five substituents, more preferably from one to three substituents, most
 10 preferably from one to two substituents, independently selected from the list of substituents.

Under standard nomenclature used throughout this disclosure, the terminal portion of the designated side chain is described first, followed by the
 15 adjacent functionality toward the point of attachment. Thus, for example, a "phenylC₁-C₆ alkylamidoC₁-C₆alkyl" substituent refers to a group of the formula



20 The compounds of the present invention may also be present in the form of a pharmaceutically acceptable salt. The pharmaceutically acceptable salt generally takes a form in which the basic nitrogen is protonated with an inorganic or organic acid. Representative organic or inorganic acids include hydrochloric, hydrobromic, hydriodic, perchloric, sulfuric, nitric, phosphoric,
 25 acetic, propionic, glycolic, lactic, succinic, maleic, fumaric, malic, tartaric, citric, benzoic, mandelic, methanesulfonic, hydroxyethanesulfonic, benzenesulfonic, oxalic, pamoic, 2-naphthalenesulfonic, p-toluenesulfonic, cyclohexanesulfamic, salicylic, saccharinic or trifluoroacetic.

30 Where the compounds according to this invention have at least one chiral center, they may accordingly exist as enantiomers. Where the compounds possess two or more chiral centers, they may additionally exist as

diastereomers. It is to be understood that all such isomers and mixtures thereof are encompassed within the scope of the present invention. Furthermore, some of the crystalline forms for the compounds may exist as polymorphs and as such are intended to be included in the present invention.

5 In addition, some of the compounds may form solvates with water (i.e., hydrates) or common organic solvents, and such solvates are also intended to be encompassed within the scope of this invention.

The term "subject" as used herein, refers to an animal, preferably a mammal, most preferably a human, who has been the object of treatment, observation or experiment.

10

The term "therapeutically effective amount" as used herein, means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes alleviation of the symptoms of the disease or disorder being treated.

15

As used herein, unless otherwise noted alkyl and alkoxy whether used alone or as part of a substituent group, include straight and branched chains having 1 to 8 carbon atoms, or any number within this range. For example, alkyl radicals include methyl, ethyl, propyl, isopropyl, *n*-butyl, isobutyl, *sec*-butyl, *t*-butyl, *n*-pentyl, 3-(2-methyl)butyl, 2-pentyl, 2-methylbutyl, neopentyl, *n*-hexyl, 2-hexyl and 2-methylpentyl. Alkoxy radicals are oxygen ethers formed from the previously described straight or branched chain alkyl groups. Cycloalkyl groups contain 3 to 8 ring carbons and preferably 5 to 7 carbons. Similarly, alkenyl and alkynyl groups include straight and branched chain alkenes and alkynes having 1 to 8 carbon atoms, or any number within this range.

20

25

30

The term "aryl" as used herein refers to an unsubstituted or substituted aromatic group such as phenyl and naphthyl. The term "aroyl" refers to the group -C(O)-aryl.

The term "heteroalkyl" as used herein represents an unsubstituted or substituted stable three to seven membered monocyclic saturated ring system which consists of carbon atoms and from one to three heteroatoms selected from N, O or S, and wherein the nitrogen or sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized. The heteroalkyl group may be attached at any heteroatom or carbon atom which results in the creation of a stable structure. Examples of such heteroalkyl groups include, but are not limited to azetidiny, piperidiny, pyrrolidiny, piperaziny, oxopiperaziny, oxopiperidiny, oxoazepiny, azepiny, tetrahydrofurany, dioxolany, tetrahydroimidazoly, tetrahydrothiazoly, tetrahydrooxazoly, tetrahydropyrany, morpholiny, thiomorpholiny, thiamorpholiny sulfoxide, thiamorpholiny sulfone and oxadiazoly. Preferred heteroalkyl groups include pyrrolidiny, piperidiny, piperaziny, morpholiny, azetidiny and tetrahydrothiazoly.

The term "heteroaryl" as used herein represents an unsubstituted or substituted stable five or six membered monocyclic aromatic ring system or an unsubstituted or substituted nine or ten membered benzo-fused heteroaromatic ring system or bicyclic heteroaromatic ring system which consists of carbon atoms and from one to four heteroatoms selected from N, O or S, and wherein the nitrogen or sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized. The heteroaryl group may be attached at any heteroatom or carbon atom that results in the creation of a stable structure. Examples of heteroaryl groups include, but are not limited to pyridyl, pyridaziny, thienyl, furany, imidazoly, isoxazoly, oxazoly, pyrazoly, pyrroly, thiazoly, thiadiazoly, triazoly, benzimidazoly, benzofurany, benzothienyl, benzisoxazoly, benzoxazoly, benzopyrazoly, indoly, benzothiazoly, benzothiadiazoly, benzotriazoly adeniny or quinoliny. Preferred heteroaryl groups include pyridyl, pyrroly, pyraziny, thiadiazoly, pyrazoly, thienyl, triazoly and quinoliny.

The term "aralkyl" means an alkyl group substituted with one, two or three aryl groups (e.g., benzyl, phenylethyl, diphenylmethyl, triphenylmethyl). Similarly, the term "aralkoxy" indicates an alkoxy group substituted with an aryl group (e.g., benzyloxy). The term aminoalkyl refers to an alkyl group substituted with an amino group (*i.e.*, -alkyl-NH₂). The term "alkylamino" refers to an amino group substituted with an alkyl group (*i.e.*, -NH-alkyl). The term "dialkylamino" refers to an amino group which is disubstituted with alkyl groups wherein the alkyl groups can be the same or different (*i.e.*, -N-[alkyl]₂).


The term "acyl" as used herein means an organic radical having 1 to 6 carbon atoms (branched or straight chain) derived from an organic acid by removal of the hydroxyl group.

The term "oxo" refers to the group =O.


The term "carbonyl" refers to the group C(O).

The term "halogen" shall include iodine, bromine, chlorine and fluorine.

The term "N(CH₂)₄" as used herein (e.g., in the Tables), refers to a pyrrolidinyl group having the structure



Similarly, "N(CH₂)₅", "N(CH₂)₃", "NCH₂S(CH₂)₂" and "N(CH₂)₂O(CH₂)₂" refer to



Similarly "C₆H₁₁" and "C₅H₉" (or "c-C₆H₁₁" and "c-C₅H₉") refer to cyclohexyl and cyclopentyl groups, respectively.

Whenever the term "alkyl" or "aryl" or either of their prefix roots appear in a name of a substituent (e.g., aralkyl, dialkylamino) it shall be interpreted as including those limitations given above for "alkyl" and "aryl." Designated numbers of carbon atoms (e.g., C₁-C₆) shall refer independently to the number of carbon atoms in an alkyl or cycloalkyl moiety or to the alkyl portion of a larger substituent in which alkyl appears as its prefix root.

As used herein, the term "phosgene equivalent" represents the class of carbonic acid derivatives which include 4-nitrophenyl chloroformate, phosgene or "COCl₂," phenyl chloroformate, triphosgene or "(CCl₃O)₂CO," carbonyldiimidazole, diethyl carbonate or diphenyl carbonate.


It is intended that the definition of any substituent or variable at a particular location in a molecule be independent of its definitions elsewhere in that molecule. It is understood that substituents and substitution patterns on the compounds of this invention can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art as well as those methods set forth herein.

20


As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combinations of the specified ingredients in the specified amounts. Accordingly, pharmaceutical compositions containing the compounds of the present invention as the active ingredient as well as methods of preparing the instant compounds are also part of the present invention.

Particularly preferred compounds of the present invention and their biological data are shown in Tables 1 and 2, following; the amino acids bear the "L" absolute configuration unless denoted otherwise. The Tables contain IC₅₀ values (μM) of the compounds in a thrombin receptor binding assay, and IC₅₀ values (μM) against platelet aggregation stimulated by thrombin.

PAGE INTENTIONALLY LEFT BLANK

Table 1. 6-Substituted Benzimidazolone Peptidomimetics

Compd	R ₁	R ₄	A ₁	A ₂ NR ₂ R ₃	IC ₅₀ (μM)	
					Thr GFP Aggr ^a	Thr Recptr Bdg ^b
1	CH ₂ N(CH ₂) ₄	2,6-DiCl-Ph ^c	3,4-DiF-Phe ^d	Dbu ^e -NHBn	0.39	1.6
4	CH ₂ NMe ₂	2-Me-Ph	3,4-DiF-Phe	Dbu-NHBn	0.23	0.7
5	CH ₂ N(CH ₂) ₄	2-Me-Ph	3,4-DiF-Phe	Dbu-NHBn	0.28	0.4
6	CH ₂ N(CH ₂) ₄	4-F-Ph	3,4-DiF-Phe	Dbu-NHBn	6	0.6
8	CH ₂ N(CH ₂) ₄	4-F-Ph	3,4-DiF-Phe	4-PyrAla ^f -NH(CH ₂) ₂ NH ₂	5.8	0.5

Table 2. 5-Substituted Benzimidazolone Peptidomimetics

Compd	R ₁	R ₄	A ₁	A ₂ NR ₂ R ₃	IC ₅₀ (μM)	
					Thr GFP Aggr ^a	Thr Recptr Bdg ^b
2	CH ₂ N(CH ₂) ₄	2-Me-Ph	3,4-DiF-Phe	Dbu-NHBn	7	3
3	CH ₂ NMe ₂	2-Me-Ph	3,4-DiF-Phe	Dbu-NHBn	7.7	1.9
7	CH ₂ N(CH ₂) ₄	2,6-DiCl-Ph	3,4-DiF-Phe	Dbu-NHBn	5.5	1.7

^a Thrombin-induced gel-filtered platelet aggregation assay.

^b Thrombin receptor (PAR-1) binding assay. ^c 2,6-Dichlorophenyl.

^d 3,4-Difluorophenylalanine. ^e 2,4-Diaminobutyric acid. ^f 4-Pyridylalanine.


The antagonists of the present invention may be prepared via a convergent solution-phase synthesis by coupling an aminobenzimidazolone intermediate **AAG4** with a dipeptide amine **AAG6** via a urea linkage as described in the general Scheme AAGeneric. The appropriately nitro substituted benzimidazolone **AAG1** (Scheme AAGeneric) was alkylated with a substituted aralkyl or heteroaryl alkyl halide and a base such as sodium hydride in a dipolar aprotic solvent such as DMF to give **AAG2** as a mixture of two regioisomers. Two isomers were separated by silica gel column and then alkylated, with an aminoalkyl halide and a base such as sodium hydride in a dipolar aprotic solvent such as DMF to give two regioisomers **AAG3**, respectively. Reduction of nitro group in **AAG3** in a classical manner with for example iron and acetic acid or with a newer method such as dimethyl hydrazine and iron to gave aminobenzimidazolone intermediate **AAG4**.

Dipeptide amine **AAG6** can be synthesized from the corresponding protected amino acids using standard peptide coupling conditions. Thus, an Fmoc protected amino-acid (A_2) **AAG5** (Scheme AAGeneric) was coupled to amine R_2R_3NH using a coupling agent, such as dicyclohexylcarbodiimide (DCC) or diisopropylcarbodiimide (DIC) and 1-hydroxybenzotriazole (HOBT) in a dipolar aprotic solvent such as DMF to give the amide. The amide was then Fmoc deprotected with a dialkylamine in a dipolar aprotic solvent, such as diethylamine in acetonitrile. The resulting amine was coupled to the second Fmoc protected amino-acid (A_1) in the same way with a coupling agent, such as DIC and HOBT in a dipolar aprotic solvent such as DMF to give the dipeptide. The dipeptide was then Fmoc deprotected as above with a dialkylamine in a dipolar aprotic solvent such as acetonitrile to afford dipeptide amine **AAG6**.

Aminobenzimidazolone intermediate **AAG4** was treated with a phosgene equivalent such as 4-nitrophenyl chloroformate, phosgene or " $COCl_2$," phenyl chloroformate, triphosgene or " $(CCl_3O)_2CO$," carbonyldiimidazole, diethyl carbonate or diphenyl carbonate and a base such as diisopropylethylamine in a solvent such as dichloromethane, and to this was then added dipeptide

amine **AAG6** to give a urea. Removal of the protecting group, if necessary, such as the Boc group with an acid such as trifluoroacetic acid from the side chain of dipeptide afforded final targets **AAG7**.





SCHEME AAGeneric




As a typical example of this convergent solution-phase method, Compound 1 may be synthesized as described in Scheme AA. Thus, treatment of 5-nitrobenzimidazolone **AA1** with 2,6-diCl-Bn-Br in the presence of NaH in DMF afforded two regioisomers **AA2a** and **AA3a** (ca. 1:1 ratio),
5 which were then separated by flash column. A small amount of bis-alkylated product was also obtained. The isolated **AA3a** was further alkylated with 2-chloroethylpyrrolidine by using NaH as a base in DMF to give the di-alkylated product, which was then subjected to nitro reduction with $\text{Me}_2\text{NNH}_2/\text{FeCl}_3$ to provide aminobenzimidazolone intermediate **AA4**. Coupling of N- α -Fmoc-N- γ -
10 Boc-diaminobutyric acid (**AA5**) with benzyl amine in the presence of DCC and HOBT was followed by de-protection of the Fmoc group with diethylamine. The resulting intermediate was coupled with Fmoc-3,4-diF-Phe-OH using DIC/HOBT and treated with diethylamine to give dipeptide amine **AA6**. Urea formation
15 **AA4** in the presence of 4-nitrophenylchloroformate was followed by de-protection of the Boc group with TFA to afford the target compound 1.

In an alternative approach to the targets, the preferred intermediate **AA3** could be prepared via a regioselective method. For example, reductive
20 alkylation of 4-nitro-1,2-phenylenediamine **AB1** with one equivalent of 4-fluorobenzaldehyde in the presence of sodium triacetoxyborohydride gave mono-alkylated product **AB2** which was then cyclized with N,N'-disuccinimidyl carbonate in acetonitrile to give **AA3c** (Scheme AB). By following the same
25 procedure as described in Scheme AA, the target 6 was obtained from intermediate **AA3c**.

SCHEME AA



SCHEME AB



The side-chain amine in antagonists such as Compound 1 and Compound 8 may be converted to other functional groups such as acetamidine and guanidine by using standard procedures. For example, the acetamidine and guanidine groups can be introduced by treating the side-chain amine with
5 S-2-naphthylmethyl thioacetimidate hydrobromide and 2-methyl-2-thiopseudourea, respectively.

The thioureidoindoles [when X is S, as in general formula (I)] may be prepared as described hereinafter. The aminobenzimidazolone substrate is
10 reacted with thiocarbonyldiimidazole in a chlorinated solvent and then with the imidazole by-product filtered from the solution. The solution can then be concentrated to afford the N-imidazolyl-N'-benzimidazolonyl-thioamide. This intermediate is then reacted with a peptide amine in a polar, aprotic solvent with heating (from about 80 °C to about 100 °C) to afford the N-peptido-N'-
15 benzimidazolonyl-thiourea product.

Amidobenzimidazolone targets [when p is 0 and X is O, as in general formula (I)] may be prepared from a dipeptide amine **AAG6** (Scheme AAGeneric) and a benzimidazolone carboxylic acid intermediate by using
20 standard coupling conditions such as DCC/HOBt. The required benzimidazolone carboxylic acid intermediates can be prepared by using the method as described for aminobenzimidazolone intermediate **AAG4** in Scheme AAGeneric from the appropriate benzimidazolone carboxylic acid esters.

25 The utility of the compounds to treat PAR-1 mediated disorders (e.g., thrombotic disorders) can be determined according to the procedures described herein. The present invention therefore provides a method of treating PAR-1 mediated disorders (e.g., thrombotic disorders) in a subject in need thereof which comprises administering any of the compounds as defined
30 herein in a quantity effective to treat PAR-1 mediated disorders. The compound may be administered to a patient by any conventional route of administration, including, but not limited to, intravenous, oral, subcutaneous, intramuscular, intradermal and parenteral.

The present invention also provides pharmaceutical compositions comprising one or more compounds of this invention in association with a pharmaceutically acceptable carrier.

5

To prepare the pharmaceutical compositions of this invention, one or more compounds of formula (I) or salt thereof of the invention as the active ingredient, is intimately admixed with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques, which carrier may take
10 a wide variety of forms depending on the form of preparation desired for administration, e.g., oral or parenteral such as intramuscular. In preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed. Thus, for liquid oral preparations, such as, for example, suspensions, elixirs and solutions, suitable carriers and additives include
15 water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like; for solid oral preparations such as, for example, powders, capsules, caplets, gelcaps and tablets, suitable carriers and additives include starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form,
20 in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be sugar coated or enteric coated by standard techniques. For parenterals, the carrier will usually comprise sterile water, though other ingredients, for example, for purposes such as aiding solubility or for preservation, may be included. Injectable suspensions may also be prepared, in which case appropriate liquid carriers, suspending agents and the like may be employed. The pharmaceutical compositions herein will contain, per dosage unit, e.g., tablet, capsule, powder, injection, teaspoonful and the like, an amount of the active ingredient necessary to deliver an effective dose as
25 described above. The pharmaceutical compositions herein will contain, per unit dosage unit, e.g., tablet, capsule, powder, injection, suppository, teaspoonful and the like, from about 0.03 mg/kg to about 100 mg/kg (preferred from about 0.1 mg/kg to about 30 mg/kg) of a compound of the present invention and may be given at a dosage from about 0.1 mg/kg/day to about
30 300 mg/kg/day (preferred from about 1 mg/kg/day to about 50 mg/kg/day). The dosages, however, may be varied depending upon the requirement of the patients, the severity of the condition being treated and the compound being

employed. The use of either daily administration or post-periodic dosing may be employed.

Preferably these compositions are in unit dosage forms such as tablets, pills, capsules, powders, granules, sterile parenteral solutions or suspensions, metered aerosol or liquid sprays, drops, ampoules, autoinjector devices or suppositories for oral parenteral, intranasal, sublingual or rectal administration, or for administration by inhalation or insufflation. Alternatively, the composition may be presented in a form suitable for once-weekly or once-monthly administration; for example, an insoluble salt of the active compound, such as the decanoate salt, may be adapted to provide a depot preparation for intramuscular injection. For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention or a pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type described above containing from about 0.1 mg to about 500 mg of the active ingredient of the present invention. The tablets or pills of the novel composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of material can be used for such enteric layers or coatings, such

materials including a number of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions, include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinyl-pyrrolidone or gelatin.

Where the processes for the preparation of the compounds according to the invention give rise to mixture of stereoisomers, these isomers may be separated by conventional techniques such as preparative chromatography. The compounds may be prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The compounds may, for example, be resolved into their components enantiomers by standard techniques, such as the formation of diastereomeric pairs by salt formation with an optically active acid, such as (-)-di-p-toluoyl-d-tartaric acid and/or (+)-di-p-toluoyl-l-tartaric acid followed by fractional crystallization and regeneration of the free base. The compounds may also be resolved by formation of diastereomeric esters or amides, followed by chromatographic separation and removal of the chiral auxiliary. Alternatively, the compounds may be resolved using a chiral HPLC column.

During any of the processes for preparation of the compounds of the present invention, it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in Protective Groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, Protective Groups in Organic

Synthesis, John Wiley & Sons, 1991. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

5 The method of treating PAR-1 mediated disorders described in the present invention may also be carried out using a pharmaceutical composition comprising any of the compounds as defined herein and a pharmaceutically acceptable carrier. The pharmaceutical composition may contain between about 0.01 mg to about 100 mg, preferably from about 5 to about 50 mg, of the compound, and may be constituted into any form suitable for the mode of
10 administration selected. Carriers include necessary and inert pharmaceutical excipients, including, but not limited to, binders, suspending agents, lubricants, flavorants, sweeteners, preservatives, dyes, and coatings. Compositions suitable for oral administration include solid forms, such as pills, tablets, caplets, capsules (each including immediate release, timed release and
15 sustained release formulations), granules, and powders, and liquid forms, such as solutions, syrups, elixirs, emulsions, and suspensions. Forms useful for parenteral administration include sterile solutions, emulsions and suspensions.

Advantageously, compounds of the present invention may be
20 administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore, compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a
25 transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic
30 pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders; lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include, without limitation, starch, gelatin, natural

sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, 5 agar, bentonite, xanthan gum and the like.

The liquid forms in suitably flavored suspending or dispersing agents such as the synthetic and natural gums, for example, tragacanth, acacia, methyl cellulose and the like. For parenteral administration, sterile 10 suspensions and solutions are desired. Isotonic preparations which generally contain suitable preservatives are employed when intravenous administration is desired.

The compound of the present invention can also be administered in the 15 form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

20 Compounds of the present invention may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol, polyhydroxyethylaspartamidephenol, or polyethyleneoxidepoly 25 ne substituted with palmitoyl residue. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, 30 polydihydropyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

Compounds of this invention may be administered in any of the foregoing compositions and according to dosage regimens established in the art whenever treatment of PAR-1 mediated disorders is required.

5 The daily dosage of the products may be varied over a wide range from about 0.01 mg to about 1,000 mg per adult human per day. For oral administration, the compositions are preferably provided in the form of tablets containing about 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, 50.0, 100, 150, 200, 250 and 500 mg of the active ingredient for the symptomatic
10 adjustment of the dosage to the patient to be treated. An effective amount of the drug is ordinarily supplied at a dosage level of from about 0.01 mg/kg to about 100 mg/kg of body weight per day. Preferably, the range is from about 0.03 mg/kg to about 10 mg/kg of body weight per day. The compounds may be administered on a regimen of about 1 time per day to about 4 times per day.

15

Optimal dosages to be administered may be readily determined by those skilled in the art, and will vary with the particular compound used, the mode of administration, the strength of the preparation, the mode of administration, and the advancement of the disease condition. In addition,
20 factors associated with the particular patient being treated, including patient age, weight, diet and time of administration, will result in the need to adjust dosages.

Biology

25

The compounds of the present invention are thrombin receptor (PAR-1) antagonists. The compounds interrupt platelet activation induced by thrombin's proteolytic cleavage of its platelet surface receptor, and thereby inhibit platelet aggregation. Such compounds are, therefore, useful in treating
30 platelet-mediated thrombotic disorders (e.g., arterial and venous thrombosis, acute myocardial infarction, reocclusion following thrombolytic therapy and angioplasty, and a variety of vaso-occlusive disorders) and other PAR-1 mediated disorders.

***In Vitro* Thrombin Receptor Binding Assay**

CHRF membranes (Jones, *Biochim. Biophys. Acta* 1992, 1136, 272) are
5 thawed from -70°C, centrifuged at maximum speed for 5 min, washed twice
with binding buffer (50 mM HEPES containing 5 mM MgCl₂ and 0.1% BSA),
and re-suspended in binding buffer (25 µg/100 mL). 100 µL of membranes are
added to the 24-Wallac plates and delivered to the Tomtech apparatus. In a
typical experiment, 6 µL of samples (from a 125 µg/mL intermediary plate, 20%
10 DMSO) and 44 µL buffer are delivered to the plates (final conc. of compounds
is 3.7 µg/mL, 0.6% DMSO). Similarly, 6 µL 20% DMSO and 44 µL buffer are
delivered to both column 1 (NSB) and column 12 (TB). 10 µL Ser-pFPhe-Har-
Leu-Har-Lys-Tyr-NH₂ (721-40; 500 µM in deionized water) is added to column
1. 50 µL tritiated 721-40 (specific activity 46 Ci/mmol) is added to all the wells.
15 The plates are mixed well for 20 seconds, incubated for 30 min, and then
harvested with 10 mM HEPES/138 mM NaCl using the Skatron harvester. The
filters (GF/C Brandel FPXLR 296) are presoaked 3 h in 0.5% polyethylenimine
in HEPES/0.1M N-acetylglucosamine) are set in saran wrap and dried for 3
min in the microwave, and placed in sample bags (Wallac 1450-432). 4.5 mL
20 scintillation fluid (Wallac, Betaplate Scint 1205-440) is added. The bags are
sealed, placed in filter cassettes (Wallac 1450-104), and analyzed on the
microbeta counter.

***In Vitro* Inhibition Of Thrombin-Induced Gel-Filtered Platelet Aggregation Assay**

The percentage of platelet aggregation is calculated as an increase in
light transmission of compound-treated platelet concentrate vs. control-treated
platelet concentrate. Human blood is obtained from drug free, normal donors
30 in tubes containing 0.13M sodium citrate. Platelet rich plasma (PRP) is
collected by centrifugation of whole blood at 200 x g for 10 min at 25°C. The
PRP (5 mL) is gel filtered through Sepharose 2B (bed volume 50 mL), and the
platelet count is adjusted to 2x10⁷ platelets per sample. The following
constituents are added to a siliconized cuvette: concentrated platelet filtrate
35 and Tyrode's buffer (0.14M NaCl, 0.0027M KCl, 0.012M NaHCO₃, 0.76 mM
Na₂HPO₄, 0.0055M glucose, 2 mg/mL BSA and 5.0 mM HEPES @ pH 7.4) in
an amount equal to 350 µL, 50 µL of 20 mM calcium and 50 µL of the test

compound. Aggregation is monitored in a BIODATA aggregometer for the 3 min following the addition of agonist (thrombin 50 μ L of 1 unit/mL).

5 Tables 1 and 2 show the biological activity of the compounds of the present invention. Tables 1 and 2 contain IC_{50} values (μ M) of the compounds against platelet aggregation stimulated by thrombin and IC_{50} values (μ M) in a thrombin receptor (PAR-1) binding assay.

Examples

10 **General Procedures:** Protected amino acids were purchased from Novabiochem, Bachem Bioscience, Advanced ChemTech or SyntheTech. All other chemicals were obtained from commercial suppliers and used without further purification. 1H and ^{13}C NMR spectra were recorded on a Bruker AC
15 300B (300 MHz proton) or a Bruker AM-400 (400 MHz proton) spectrometer with Me_4Si as an internal standard (s = singlet, d = doublet, t = triplet, br = broad). APCI-MS and ES-MS were recorded on a VG Platform II mass spectrometer; methane was used for chemical ionization, unless noted otherwise. Accurate mass measurements were obtained by using a VG ZAB
20 2-SE spectrometer in the FAB mode. TLC was performed with Whatman 250- μ m silica gel plates. Preparative TLC was performed with Analtech 1000- μ m silica gel GF plates. Flash column chromatography was conducted with flash column silica gel (40-63 μ m) and column chromatography was conducted with standard silica gel. HPLC separations were carried out on three Waters
25 PrepPak® Cartridges (25 x 100 mm, Bondapak® C18, 15-20 μ m, 125 Å) connected in series; detection was at 254 nm on a Waters 486 UV detector. Analytical HPLC was carried out on a Supelcosil ABZ+PLUS column (5 cm x 2.1 mm), with detection at 254 nm on a Hewlett Packard 1100 UV detector. Microanalysis was performed by Robertson Microlit Laboratories, Inc.

30

In the examples and throughout this application, the following abbreviations have the meanings recited hereinafter:

	Ac	Acetyl
	ACN	Acetonitrile
	Bn	Benzyl
	Boc	t-Butoxycarbonyl
5	DCC	1,3-Dicyclohexylcarbodiimide
	DCM	Dichloromethane
	DIC	Diisopropylcarbodiimide
	DIEA	Diisopropylethylamine
	DMAP	4-Dimethylaminopyridine
10	DMF	N, N-Dimethylformamide
	Et	Ethyl
	Fmoc	9-Fluorenylmethoxycarbonyl
	h	Hour
15	HBTU	2-(1H-Benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
	HOAc	Acetic acid
	HOBT	Hydroxybenzotriazole
	Me	Methyl
	min	Minute
20	rt	room temperature
	THF	Tetrahydrofuran
	TFA	Trifluoroacetic acid

Example 1

25 Synthesis of Compound 1 (Scheme AA)

30 Benzenepropanamide, *N*-[(1*S*)-3-amino-1-
 [[[phenylmethyl]amino]carbonyl]propyl]- α -[[[3-(2,6-dichlorophenyl)methyl]-2,3-
 dihydro-2-oxo-1-[2-(1-pyrrolidinyl)ethyl]-1*H*-benzimidazol-5-
 yl]amino]carbonyl]amino]-3,4-difluoro-, (α *S*)- (1)

35 5-Nitroimidazolone **AA1** (1.44 g, 8.04 mm) in dry DMF (20 mL) under
 argon was cooled to about 0°C and 75% NaH oil dispersion (0.284 g, 8.86
 mm) was added portionwise and stirred for about 5 min.
 2,6-Dichlorobenzylbromide (1.92 g, 8.0 mm) dissolved in dry DMF (20 mL) was
 added over about 5 min and the reaction was stirred at about 0°C for about 30
 min and then at about rt for about 3 h. The solution was added slowly to

vigorously stirring water (700 mL) and after about 1 h, a white solid was filtered (2.38 g) that contained a mixture of **AA2a** and **AA3a** (ca. 1:1 ratio). A portion (1.5 g) was purified by flash column chromatography using DCM:ethyl acetate (2:1) to give pure **AA2a** and **AA3a**. A sample of **AA3a** (105 mg, 0.30 mm) was dissolved in dry DMF (6 mL), 75% NaH (30 mg, 0.94 mm) was added and the mixture was stirred for about several min at about rt. 2-chloroethylpyrrolidine hydrochloride (103 mg, 0.60 mm) was added and the mixture was stirred at about 50°C for about 2 h. The reaction was cooled and partitioned between chloroform:2-propanol (10:1) and water; the organic solution was washed twice with water, once with brine, dried (K_2CO_3) and evaporated *in vacuo* to a light yellow solid (130 mg). This solid was dissolved in MeOH (8.0 mL) and ferric chloride hexahydrate (30 mg, 0.11 mm), charcoal (60 mg, 5 mm) and 1,1 dimethyl hydrazine (1.5 g, 25 mm) were added and the reaction was refluxed for about 3 h. The solution was cooled to about rt, filtered through dicalite and the filtrate was evaporated *in vacuo* to a clear oil. The oil was dissolved in 1N HCl (25 mL) and washed twice with ethyl ether. The acid was made alkaline with dilute NaOH and extracted with DCM, which was washed twice with sodium bicarbonate solution, once with brine, dried (K_2CO_3) and evaporated *in vacuo* to give solid **AA4** (100 mg). Fmoc- α -N-Boc- γ -N-diaminobutyric acid **AA5** (10.8 g, 24.5 mmol) was stirred in ACN (300 mL) under argon as HOBT (3.75 g, 24.4 mmol) was added, followed by benzylamine (2.6 g, 24.3 mmol). DCC (10.4 g, 48.7 mmol) was also then added and the reaction was stirred at about rt for about 3 h, whereupon a white solid was filtered and washed with cold ACN (14.4 g). The solid was stirred in ACN (500 mL) containing diethyl amine (25 mL) for about 2 h and a small quantity of solid was filtered; the filtrate was evaporated *in vacuo* to an oil, which was triturated three times with hexane (400 mL each) to white solid. The solid was dissolved in ACN (400 mL), then HOBT (2.9 g, 19.1 mmol) and Fmoc-3,4 difluorophenylalanine (8.1 g, 19.1 mmol) were added, followed by DIC (4.81 g, 38.2 mmol) and stirred at about rt for about 16 h. The reaction was cooled in an ice bath and a white solid was filtered then washed with cold ACN. The solid was stirred in ACN (350 mL) containing diethylamine (35 mL) for about 5 h and evaporated *in vacuo* to a white solid. The solid was triturated three times with hexane, dissolved in chloroform (250 mL), dried (Na_2SO_4) and evaporated *in vacuo* to a white solid **AA6** (8.0 g). 4-Nitrophenyl chloroformate (20 mg, 0.10 mm) in DCM (5 mL) was cooled to about -20°C and **AA4** (40 mg, 0.10 mm) and diisopropylethylamine (13 mg, 0.10 mm) in DCM (1.5 mL) were added and stirred for about 30 min. The amine **AA6** (50 mg, 0.10 mm) and

diisopropylethylamine (13 mg, 0.10 mm) in DCM (2.0 mL) were added to the solution at about -20°C and stirred at about -20°C for about 30 min. The mixture was then allowed to come to about rt, whereupon the reaction was stirred for about another 16 h. A light yellow solid was filtered (45 mg) and then stirred into DCM (7 mL) containing anisole (10 mg) as TFA (3.0 mL) was added under argon. The reaction was stirred at about rt for about 1.5 h and then evaporated *in vacuo* to an oil, which was triturated three times with ethyl ether to give Compound 1, a white solid, as its trifluoroacetate salt. ¹H NMR (CD₃OD) δ 1.9 - 2.2 (m, 6H), 2.9 - 3.1 (m, 3H), 3.1 - 3.3 (m, 4H), 3.6 (dd, 2H), 3.8 (m, 2H), 4.25 (dd, 2H), 4.35 (s, 2H), 4.45 (dd, 2H), 5.32 (s, 2H), 7.0 - 7.5 (m, aromatics). ES-MS *m/z* 821 (MH⁺). FAB-HRMS calcd for C₄₁H₄₄Cl₂F₂N₈O₄ + H⁺: 821.2909, found 821.2918.

Example 2

Synthesis of Compound 6 (Scheme AB)

Benzenepropanamide, *N*-[(1*S*)-3-amino-1-
[[[(phenylmethyl)amino]carbonyl]propyl]-α-[[[3-(4-fluorophenyl)methyl]-2,3-
dihydro-2-oxo-1-[2-(1-pyrrolidinyl)ethyl]-1*H*-benzimidazol-5-
yl]amino]carbonyl]amino]-3,4-difluoro-, (α*S*)- (6)

4-Nitro-1,2-phenylene diamine (**AB1**, 2.00 g, 13.0 mmol) was suspended in DCE (100 mL) followed by the addition of 4-fluorobenzaldehyde (1.39 mL, 13.0 mmol), acetic acid (0.74 mL, 13.0 mmol), and sodium triacetoxyborohydride (5.51 g, 26.0 mmol). Reaction was stirred at ambient temperature for 24 h. Additional sodium triacetoxyborohydride (5.51 g, 26.0 mmol) was added and the reaction was stirred at ambient temperature for 24 h. The reaction mixture was diluted with EtOAc (175 mL), washed with H₂O (2X50 mL), brine (50 mL), dried (Na₂SO₄), and evaporated *in vacuo* to give a solid. The crude product was purified by column chromatography (silica gel, 20% EtOAc/80% Hexane to 40% EtOAc/60% Hexane) to afford **AB2** (1.42 g, 42%). To a solution of **AB2** (1.39 g, 5.3 mmol) in acetonitrile (80 mL) was added *N,N'*-disuccinimidyl carbamate (1.36 g, 5.3 mmol) in acetonitrile (80 mL) and the mixture was stirred at 75 °C. After 24 h, additional *N,N'*-disuccinimidyl carbonate (1.36 g, 5.3 mmol) was added and the reaction mixture stirred at 75 °C for another 24 h. The reaction was still not complete and additional *N,N'*-disuccinimidyl carbonate (1.36 g, 5.3 mmol) was added. The stirring was continued at 75 °C for

4 h, and then the reaction mixture was evaporated *in vacuo* to give a solid. The crude product was purified by column chromatography (silica gel-30% EtOAc/70% Hexane to 50% EtOAc/50% Hexane) to afford **AA3c** (0.95 g, 62%). ¹H NMR (DMSO) δ 5.13 (s, CH₂), 7.16-7.21 (m, 3H, H-11, H-13, and H-3), 7.39-7.44 (m, 2H, H-10 and H-13), 7.97-8.01 (m, 2H, H-4 and H-5), 11.76 (s, NH). To a solution of **AA3c** (0.50 g, 1.7 mmol) in dry DMF (24 ml) was added 60 % NaH (0.20 g, 5.1 mmol) and the mixture was stirred at rt for several min. 2-Chloroethylpyrrolidine hydrochloride (0.57g, 3.4 mmol) was added and the mixture was stirred at about 50 °C overnight. The reaction was cooled to rt and additional 60% NaH (32.0 mg, 0.85 mmol) and 2-chloroethylpyrrolidine hydrochloride (57.8 mg, 0.34 mmol) was added and the mixture was stirred at about 50 °C for 5 h. The reaction was cooled and partitioned between chloroform:2-propanol (10:1) and water; the organic solution was washed twice with water, twice with brine, dried (K₂CO₃) and evaporated *in vacuo* to give a light yellow solid (0.72 g). The solid (0.70 g, 1.8 mmol) was dissolved in MeOH (50 mL) and ferric chloride hexahydrate (0.19 g, 0.70 mmol), charcoal, (0.39 g) and 1,1-dimethyl hydrazine (12.2 mL, 160 mmol) were added and the reaction was refluxed for 2 h. The solution was cooled to rt, filtered through celite and the filtrate was evaporated *in vacuo* to give an oil. The oil was dissolved in 1N HCl (150 mL) and washed twice with ether. The acid was basified with aqueous NaOH, extracted with DCM, which was washed twice with sodium bicarbonate solution, twice with brine, dried (K₂CO₂) and evaporated *in vacuo* to give solid **AB3** (0.39 g). 4-Nitrophenyl chloroformate (0.028 g, 0.14 mmol) in DCM (5 mL) was cooled to about -20 °C and **AB3** (50 mg, 0.14 mmol) and diisopropylethylamine (18 mg, 0.14 mmol) in DCM (1.5 mL) were added and stirred for 2 h. The amine **AA6** (0.079 g, 0.14 mmol) and diisopropylethylamine (18 mg, 0.14 mmol) in DCM (2.0 mL) were added to the solution at about -20 °C and stirred at about -20 °C for 1h. The mixture was then allowed to warm up to rt, whereupon the reaction was stirred for about 18 h. A light yellow solid was collected by filtration and then treated with 30% TFA/DCM/1% anisole (8 mL) under N₂. The reaction was stirred at rt for 1 h and was then blown down with N₂ to an oil, which was triturated four times with ethyl ether to give compound **6**, a white solid, as its trifluoroacetate salt. ¹H NMR (CD₃OD) δ 1.95-2.20 (m, 6H), 2.96-3.61 (m, 11H), 4.29-4.33 (m, 4H), 4.47 (dd, 2H), 5.01 (d, 2H), 6.96-7.36 (m, aromatics). ES-MS *m/z* 771 (MH⁺).


Example 3

As a specific embodiment of an oral composition, 100 mg of the Compound 1 of Example 1 is formulated with sufficient finely divided lactose to
5 provide a total amount of about 580 mg to about 590 mg to fill a size O hard gel capsule.

While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be
10 understood that the practice of the invention encompasses all of the usual variations, adaptations and/or modifications as come within the scope of the following claims and their equivalents.

WHAT IS CLAIMED IS:

1. A compound of the following formula (I):



(I)

wherein:

A_1 and A_2 are each independently a D- or L-amino acid selected from the group consisting of alanine, β -alanine, arginine, homoarginine, cyclohexylalanine, citrulline, cysteine (optionally substituted with C_1 - C_4 alkyl, aryl, or arC_1 - C_4 alkyl), 2,4-diaminobutyric acid (optionally substituted with acyl, C_1 - C_4 alkyl, aroyl, amidino, or $MeC(NH)-$), 2,3 diaminopropionic acid (optionally substituted with acyl, C_1 - C_4 alkyl, aroyl, amidino, or $MeC(NH)-$), glutamine, glycine, indanylglycine, lysine (optionally substituted with acyl, C_1 - C_4 alkyl, aroyl, $MeC(NH)-$), valine, methionine, proline, serine (optionally substituted with C_1 - C_4 alkyl, aryl, or arC_1 - C_4 alkyl), homoserine (optionally substituted with C_1 - C_4 alkyl, aryl, or arC_1 - C_4 alkyl), tetrahydroisoquinoline-3-COOH, threonine (optionally substituted with C_1 - C_4 alkyl, aryl, or arC_1 - C_4 alkyl), ornithine (optionally substituted with acyl, C_1 - C_4 alkyl, aroyl, $MeC(NH)-$), and an unsubstituted or substituted aromatic amino acid selected from the group consisting of phenylalanine, heteroarylalanine, naphthylalanine, homophenylalanine, histidine, tryptophan, tyrosine, arylglycine, heteroarylglycine, aryl- β -alanine, and heteroaryl- β -alanine wherein the substituents on the aromatic amino acid are independently selected from one or more of halogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, hydroxy, C_1 - C_4 alkoxycarbonyl,

amino, amidino, guanidino, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl, heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or nitro;

- 5 R₁ is selected from amino, C₁-C₈ alkylamino, C₁-C₈ dialkylamino, arylamino, arC₁-C₈ alkylamino, C₃-C₈ cycloalkylamino, heteroalkylC₁-C₈ alkylamino, heteroalkylC₁-C₈ alkyl-N-methylamino, C₁-C₈ dialkylaminoC₁-C₈ alkylamino, -N(C₁-C₈alkyl)-C₁-C₈ alkyl-N(C₁-C₈alkyl)₂, N(C₁-C₈ alkyl)(C₁-C₈ alkenyl), -N(C₁-C₈alkyl)(C₃-C₈cycloalkyl), heteroalkyl or substituted heteroalkyl
10 wherein the substituent on the heteroalkyl is selected from oxo, amino, C₁-C₈ alkoxyC₁-C₈ alkyl, C₁-C₈ alkylamino or C₁-C₈ dialkylamino;

- R₂ and R₃ are each independently selected from hydrogen, C₁-C₈ alkyl, C₃-C₈ cycloalkyl, C₃-C₈ cycloalkylC₁-C₈ alkyl, aryl, heteroalkyl, substituted
15 heteroalkyl (wherein the substituent on the heteroalkyl is one or more substituents independently selected from C₁-C₈ alkoxycarbonyl, C₁-C₈ alkyl, or C₁-C₄ alkylcarbonyl), heteroalkylC₁-C₈ alkyl, indanyl, acetamidinoC₁-C₈ alkyl, aminoC₁-C₈ alkyl, C₁-C₈ alkylaminoC₁-C₈ alkyl, C₁-C₈ dialkylaminoC₁-C₈ alkyl, unsubstituted or substituted heteroarylC₁-C₈ alkyl or unsubstituted or
20 substituted arC₁-C₈ alkyl, wherein the substituent on the aralkyl or heteroarylalkyl group is one or more substituents independently selected from halogen, nitro, amino, C₁-C₈ alkyl, C₁-C₈ alkoxy, hydroxy, cyano, C₁-C₄ alkylcarbonyl, C₁-C₈ alkoxycarbonyl, hydroxyC₁-C₈ alkyl or aminosulfonyl; or

- 25 R₂ and R₃, together with the nitrogen to which they are attached, alternatively form an unsubstituted or substituted heteroalkyl group selected from piperidinyl, piperazinyl, morpholinyl or pyrrolidinyl, wherein the substituent is one or more substituents independently selected from C₁-C₈ alkyl C₁-C₈ alkoxycarbonyl or C₁-C₄ alkylcarbonyl;

30

- R₄ is selected from unsubstituted or substituted aryl, arC₁-C₈ alkyl, C₃-C₈ cycloalkyl, or heteroaryl, where the substituents on the aryl, arC₁-C₈ alkyl, cycloalkyl or heteroaryl group are independently selected from one or more of
35 C₁-C₄ alkylcarbonyl, C₁-C₈ alkoxycarbonyl, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl;

R₅ is selected from hydrogen or C₁-C₈ alkyl;

X is oxygen or sulfur;

- 5 m is an integer selected from 0, 1, 2 or 3;
 n is an integer selected from 2 or 3; and
 p is an integer selected from 0 or 1;

and pharmaceutically acceptable salts thereof.

10

2. The compound of Claim 1, wherein:

A₁ and A₂ are each independently an L-amino acid selected from the group consisting of alanine, β-alanine, arginine, homoarginine, cyclohexylalanine, citrulline, cysteine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), 2,4-diaminobutyric acid (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, amidino, or MeC(NH)-), 2,3 diaminopropionic acid (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, amidino, or MeC(NH)-), glutamine, glycine, indanylglycine, lysine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), valine, methionine, proline, serine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), homoserine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), tetrahydroisoquinoline-3-COOH, threonine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), ornithine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), and an unsubstituted or substituted aromatic amino acid selected from the group consisting of phenylalanine, heteroarylalanine, naphthylalanine, homophenylalanine, histidine, tryptophan, tyrosine, arylglycine, heteroarylglycine, aryl-β-alanine, and heteroaryl-β-alanine wherein the substituents on the aromatic amino acid are independently selected from one or more of halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, C₁-C₄ alkoxycarbonyl, amino, amidino, guanidino, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl, heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or nitro;

R_1 is selected from amino, C_1 - C_6 alkylamino, C_1 - C_6 dialkylamino, arylamino, arC_1 - C_6 alkylamino, heteroalkyl C_1 - C_6 alkylamino, $-N(C_1$ - C_6 alkyl)- C_1 - C_6 alkyl- $N(C_1$ - C_6 alkyl) $_2$, heteroalkyl or substituted heteroalkyl wherein the substituent on the heteroalkyl is selected from oxo, amino, C_1 - C_6 alkoxy C_1 - C_6 alkyl, C_1 - C_6 alkylamino or C_1 - C_6 dialkylamino;

R_2 is selected from hydrogen or C_1 - C_6 alkyl;

R_3 is selected from C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, C_3 - C_6 cycloalkyl C_1 - C_6 alkyl, aryl, heteroaryl C_1 - C_6 alkyl, substituted heteroaryl C_1 - C_6 alkyl wherein the substituent is C_1 - C_4 alkyl, heteroalkyl, heteroalkyl C_1 - C_6 alkyl, indanyl, acetamidino C_1 - C_6 alkyl, amino C_1 - C_6 alkyl, C_1 - C_6 alkylamino C_1 - C_6 alkyl, C_1 - C_6 dialkylamino C_1 - C_6 alkyl, arC_1 - C_6 alkyl, substituted arC_1 - C_6 alkyl wherein the substituent on the aralkyl group is one to five substituents independently selected from halogen, nitro, amino, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_1 - C_6 alkoxycarbonyl, hydroxyalkyl or aminosulfonyl;

R_2 and R_3 , together with the nitrogen to which they are attached, alternatively form an unsubstituted or substituted heteroalkyl group selected from piperidiny, piperaziny or pyrrolidinyl, wherein the substituent is independently one or two substituents selected from C_1 - C_6 alkyl;

R_4 is selected from unsubstituted or substituted aryl, arC_1 - C_6 alkyl, C_3 - C_6 cycloalkyl or heteroaryl, where the substituents on the aryl, aralkyl, cycloalkyl or heteroaryl group are independently selected from one to three substituents selected from halogen, cyano, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 alkoxycarbonyl, fluorinated C_1 - C_4 alkyl, fluorinated C_1 - C_4 alkoxy or C_1 - C_4 alkylsulfonyl;

R_5 is hydrogen;

X is oxygen; and

p is 1;

and pharmaceutically acceptable salts thereof.

3. The compound of Claim 2, wherein:

A₁ is an L-amino acid selected from the group consisting of alanine, arginine, cyclohexylalanine, glycine, proline, tetrahydroisoquinoline-3-COOH, and an unsubstituted or substituted aromatic amino acid selected from the group consisting of phenylalanine, naphthylalanine, homophenylalanine, and
5 O-methyl tyrosine, wherein the substituents on the aromatic amino acid are independently selected from one to five of halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, C₁-C₄ alkoxycarbonyl, amino, amidino, guanidino, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl, heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or nitro;

10

A₂ is an L-amino acid selected from the group consisting of alanine, β-alanine, arginine, citrulline, cysteine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), 2,4-diaminobutyric acid (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, amidino, or MeC(NH)-), 2,3- diaminopropionic acid
15 (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, amidino, or MeC(NH)-), glutamine, glycine, lysine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), valine, methionine, serine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), homoserine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), threonine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-
20 C₄ alkyl), ornithine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), and an unsubstituted or substituted aromatic amino acid selected from the group consisting of phenylalanine, heteroarylalanine, and histidine, wherein the substituents on the aromatic amino acid are independently selected from one to five of halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, C₁-C₄
25 alkoxycarbonyl, amino, amidino, guanidino, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl, heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or nitro;

30

R₂ is selected from hydrogen or C₁-C₄ alkyl; and

m is 1 and n is 2;

and pharmaceutically acceptable salts thereof.


4. The compound of Claim 3, wherein:

A₁ is an L-amino acid selected from the group consisting of alanine, arginine, cyclohexylalanine, glycine, proline, and an unsubstituted or substituted aromatic amino acid selected from the group consisting of phenylalanine, naphthylalanine, homophenylalanine, and O-methyl tyrosine, wherein the substituents on the aromatic amino acid are independently one to two substituents selected from halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, C₁-C₄ alkoxycarbonyl, amino, amidino, guanidino, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl, heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or nitro;

A₂ is an L-amino acid selected from the group consisting of alanine, β-alanine, arginine, citrulline, cysteine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), 2,4-diaminobutyric acid (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, amidino, or MeC(NH)-), 2,3-diaminopropionic acid (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, amidino, or MeC(NH)-), glutamine, glycine, lysine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), valine, methionine, serine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), homoserine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), threonine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), ornithine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), and an unsubstituted or substituted aromatic amino acid selected from the group consisting of phenylalanine, heteroarylalanine, and histidine, wherein the substituents on the aromatic amino acid are independently one to two substituents selected from halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, C₁-C₄ alkoxycarbonyl, amino, amidino, guanidino, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl, heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or nitro;

30

R₁ is selected from dimethylamino, diethylamino, di-(*n*-propyl)amino,



R₂ is selected from hydrogen, methyl or ethyl;


- 5 R₃ is selected from 2-indanyl, phenyl, cyclohexylmethyl, cyclopentyl, pyridylmethyl, furanylmethyl, 2-(4-methyl-furanyl)methyl, thienylmethyl, diphenylmethyl, 4-imidazolylethyl, 2-(4-N-methyl)imidazolylethyl, *n*-octyl, phenyl-*n*-propyl, aminoethyl, aminopropyl, amino-*n*-pentyl, dimethylaminoethyl, 4-aminophenylsulfonylaminomethyl, acetamidineylethyl, 2-N-pyrrolidinylethyl, 10 N-ethoxycarbonylpiperidiny, unsubstituted or substituted phenylethyl or unsubstituted or substituted benzyl wherein the substituents on the phenylethyl or benzyl are independently one or two substituents selected from methyl, fluorine, chlorine, nitro, methoxy, methoxycarbonyl or hydroxymethyl; or

- 15 R₂ and R₃, together with the nitrogen to which they are attached, alternatively form a heteroalkyl group selected from piperidiny, or 4-(N-methyl)piperazinyl; and

- 20 R₄ is selected from cyclohexyl, 2-naphthyl, phenylethyl, 4-fluorophenylethyl or unsubstituted or substituted phenyl, where the substituents on the phenyl are independently selected from one to two substituents selected from fluorine, chlorine, iodine, methyl, cyano or trifluoromethyl;

- 25 and pharmaceutically acceptable salts thereof.


5. The compound of Claim 4, wherein:



30

and pharmaceutically acceptable salts thereof.

6. The compound of Claim 5, wherein:



5 and pharmaceutically acceptable salts thereof.

7. The compound of Claim 6, wherein:

- 10 A₁ is selected from 3,4-Difluorophenylalanine or 4-Chlorophenylalanine;
 A₂ is selected from 2,4-Diaminobutyric acid or 4-Pyridylalanine;
 R₂ is hydrogen; and
 R₃ is selected from benzyl or 2-aminoethyl;

and pharmaceutically acceptable salts thereof.

15

8. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound of Claim 1.

9. A pharmaceutical composition made by mixing a compound of Claim 1
 20 and a pharmaceutically acceptable carrier.

10. A process for making a pharmaceutical composition comprising mixing a compound of Claim 1 and a pharmaceutically acceptable carrier.

25 11. A method of treating a condition selected from the group consisting of thrombosis, restenosis, hypertension, heart failure, arrhythmia, myocardial infarction, glomerulonephritis, reocclusion following thrombolytic therapy, reocclusion following angioplasty, inflammation, angina, stroke, atherosclerosis, ischemic conditions, a vaso-occlusive disorder,
 30 neurodegenerative disorders, Angiogenesis related disorders and cancer in a

subject in need thereof comprising administering to the subject a therapeutically effective amount of the compound of Claim 1.

12. The method of Claim 11, wherein the therapeutically effective amount of
5 the compound is from about 0.1 mg/kg/day to about 300 mg/kg/day.

13. A method of treating a condition selected from the group consisting of thrombosis, restenosis, hypertension, heart failure, arrhythmia, myocardial infarction, glomerulonephritis, reocclusion following thrombolytic therapy,
10 reocclusion following angioplasty, inflammation, angina, stroke, atherosclerosis, ischemic conditions, a vaso-occlusive disorder, neurodegenerative disorders, Angiogenesis related disorders and cancer in a subject in need thereof comprising administering to the subject a therapeutically effective amount of the composition of Claim 8.

15

14. The method of Claim 13, wherein the therapeutically effective amount of the compound is from about 0.1 mg/kg/day to about 300 mg/kg/day.

15. A method of inhibiting platelet aggregation in a subject in need thereof
20 comprising administering to the subject a therapeutically effective amount of the compound of Claim 1.


16. The method of Claim 15, wherein the therapeutically effective amount of the compound is from about 0.1 mg/kg/day to about 300 mg/kg/day.

25

17. A method of inhibiting platelet aggregation in a subject in need thereof comprising administering to the subject a therapeutically effective amount of the composition of Claim 8.

30 18. The method of Claim 17, wherein the therapeutically effective amount of the compound is from about 0.1 mg/kg/day to about 300 mg/kg/day.

19. A method of treating a condition mediated by thrombin receptor (PAR-1) in a subject in need thereof comprising administering to the subject a therapeutically effective amount of the compound of Claim 1.
20. The method of Claim 19, wherein the therapeutically effective amount of the compound is from about 0.1 mg/kg/day to about 300 mg/kg/day.
21. A method of treating a condition mediated by thrombin receptor (PAR-1) in a subject in need thereof comprising administering to the subject a therapeutically effective amount of the composition of Claim 8.
22. The method of Claim 21, wherein the therapeutically effective amount of the compound is from about 0.1 mg/kg/day to about 300 mg/kg/day.
23. A process for preparing a compound of the formula (II):



wherein:

- 20 A_1 and A_2 are each independently a D- or L-amino acid selected from the group consisting of alanine, β -alanine, arginine, homoarginine, cyclohexylalanine, citrulline, cysteine (optionally substituted with C_1 - C_4 alkyl, aryl, or arC_1 - C_4 alkyl), 2,4-diaminobutyric acid (optionally substituted with acyl, C_1 - C_4 alkyl, aroyl, amidino, or $MeC(NH)$ -), 2,3 diaminopropionic acid (optionally substituted with acyl, C_1 - C_4 alkyl, aroyl, amidino, or $MeC(NH)$ -), glutamine, glycine, indanylglycine, lysine (optionally substituted with acyl, C_1 - C_4 alkyl, aroyl, $MeC(NH)$ -), valine, methionine, proline, serine (optionally substituted with C_1 - C_4 alkyl, aryl, or arC_1 - C_4 alkyl), homoserine (optionally substituted with C_1 - C_4

alkyl, aryl, or arC₁-C₄ alkyl), tetrahydroisoquinoline-3-COOH, threonine (optionally substituted with C₁-C₄ alkyl, aryl, or arC₁-C₄ alkyl), ornithine (optionally substituted with acyl, C₁-C₄ alkyl, aroyl, MeC(NH)-), and an unsubstituted or substituted aromatic amino acid selected from the group
 5 consisting of phenylalanine, heteroarylalanine, naphthylalanine, homophenylalanine, histidine, tryptophan, tyrosine, arylglycine, heteroarylglycine, aryl-β-alanine, and heteroaryl-β-alanine wherein the substituents on the aromatic amino acid are independently selected from one or more of halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, C₁-C₄ alkoxycarbonyl,
 10 amino, amidino, guanidino, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl, C₁-C₄ alkylcarbonyl, cyano, aryl, heteroaryl, arC₁-C₄ alkyl, C₂-C₄ alkenyl, alkynyl, or nitro;

R₁ is selected from amino, C₁-C₈ alkylamino, C₁-C₈ dialkylamino, arylamino, arC₁-C₈ alkylamino, C₃-C₈ cycloalkylamino, heteroalkylC₁-C₈ alkylamino, heteroalkylC₁-C₈ alkyl-N-methylamino, C₁-C₈ dialkylaminoC₁-C₈ alkylamino, -N(C₁-C₈alkyl)-C₁-C₈ alkyl-N(C₁-C₈alkyl)₂, N(C₁-C₈ alkyl)(C₁-C₈ alkenyl), -N(C₁-C₈alkyl)(C₃-C₈cycloalkyl), heteroalkyl or substituted heteroalkyl wherein the substituent on the heteroalkyl is selected from oxo, amino, C₁-C₈ alkoxyC₁-C₈ alkyl, C₁-C₈ alkylamino or C₁-C₈ dialkylamino;
 20

R₂ and R₃ are each independently selected from hydrogen, C₁-C₈ alkyl, C₃-C₈ cycloalkyl, C₃-C₈ cycloalkylC₁-C₈ alkyl, aryl, heteroalkyl, substituted heteroalkyl (wherein the substituent on the heteroalkyl is one or more substituents independently selected from C₁-C₈ alkoxycarbonyl, C₁-C₈ alkyl, or C₁-C₄ alkylcarbonyl), heteroalkylC₁-C₈ alkyl, indanyl, acetamidinoC₁-C₈ alkyl, aminoC₁-C₈ alkyl, C₁-C₈ alkylaminoC₁-C₈ alkyl, C₁-C₈ dialkylaminoC₁-C₈ alkyl, unsubstituted or substituted heteroarylC₁-C₈ alkyl or unsubstituted or substituted arC₁-C₈ alkyl, wherein the substituent on the aralkyl or
 25 heteroarylalkyl group is one or more substituents independently selected from halogen, nitro, amino, C₁-C₈ alkyl, C₁-C₈ alkoxy, hydroxy, cyano, C₁-C₄ alkylcarbonyl, C₁-C₈ alkoxycarbonyl, hydroxyC₁-C₈ alkyl or aminosulfonyl;
 30

R₂ and R₃, together with the nitrogen to which they are attached, alternatively form an unsubstituted or substituted heteroalkyl group selected from piperidinyl, piperazinyl, morpholinyl or pyrrolidinyl, wherein the substituent
 35

is one or more substituents independently selected from C₁-C₈ alkyl C₁-C₈ alkoxy, carbonyl or C₁-C₄ alkylcarbonyl;


5 R₄ is selected from unsubstituted or substituted aryl, arC₁-C₈ alkyl, C₃-C₈ cycloalkyl, or heteroaryl, where the substituents on the aryl, arC₁-C₈ alkyl, cycloalkyl or heteroaryl group are independently selected from one or more of halogen, nitro, amino, cyano, hydroxyalkyl, C₁-C₈ alkyl, C₁-C₈ alkoxy, hydroxy, C₁-C₄ alkylcarbonyl, C₁-C₈ alkoxy, carbonyl, fluorinated C₁-C₄ alkyl, fluorinated C₁-C₄ alkoxy, C₁-C₄ alkylsulfonyl;

10


m is 1 and n is 2;

comprising reacting a compound of the formula **AAG6**:

15



with a compound of the formula **AAG4**:



20

in the presence of a phosgene equivalent to form the compound of formula (II).

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/17751

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07K5/06 C07D235/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BIOSIS, CHEM ABS Data, EPO-Internal, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 2000 MARLOWE CHARLES ET AL: "Design, synthesis and structure-activity relationship for a series of factor Xa inhibitors containing the imidazolone nucleus as a central template" Database accession no. PREV199900423632 XP000942431 Abstract abstract & ABSTRACTS OF PAPERS AMERICAN CHEMICAL SOCIETY, vol. 219, 2000, page MEDI 5 219th National Meeting of the American Chemical Society, Parts 1 and 2; New Orleans, Louisiana, USA; August 22-26, 1999</p> <p>-/--</p>	1-23

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

26 October 2000

Date of mailing of the international search report

10/11/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kronester-Frei, A

INTERNATIONAL SEARCH REPORT

Intern 1st Application No

PCT/US 00/17751

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	ISSN: 0065-7727 ---	
Y	DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1999 ZHANG HAN-CHENG ET AL: "Novel indole-based peptidomimetics as potent thrombin receptor (PAR-1) antagonists." Database accession no. PREV199900423632 XP000942432 whole abstract abstract & ABSTRACTS OF PAPERS AMERICAN CHEMICAL SOCIETY, vol. 218, no. 1-2, 1999, page MEDI 5 218th National Meeting of the American Chemical Society, Parts 1 and 2; New Orleans, Louisiana, USA; August 22-26, 1999 ISSN: 0065-7727	1-23
Y	HOEKSTRA WILLIAM J ET AL: "Thrombin receptor (PAR-1) antagonists, heterocycle-based peptidomimetics of the SFLLR agonist motif." BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, vol. 8, no. 13, 7 July 1998 (1998-07-07), pages 1649-1654, XP002151181 ISSN: 0960-894X whole document, esp. Table 1	1-23
Y	EP 0 385 850 A (UNION PHARMA SCIENT APPL) 5 September 1990 (1990-09-05) examples, claims	1-23
A	M S BERNATOWICZ ET AL: "Development of potent thrombin receptor antagonist peptides" JOURNAL OF MEDICINAL CHEMISTRY, US, AMERICAN CHEMICAL SOCIETY. WASHINGTON, vol. 39, no. 25, 6 December 1996 (1996-12-06), pages 4879-4887, XP002095053 ISSN: 0022-2623 whole article	1-23
Y	WO 92 14750 A (UNIV CALIFORNIA ; COR THERAPEUTICS INC (US)) 3 September 1992 (1992-09-03) claims 10ff -----	1-23

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/17751

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0385850 A	05-09-1990	FR 2643903 A	07-09-1990
		AU 5061190 A	06-09-1990
		CA 2011222 A	03-09-1990
		JP 3014579 A	23-01-1991
		NZ 232621 A	26-02-1991
		PT 93328 A	07-11-1990
		ZA 9001602 A	28-12-1990

WO 9214750 A	03-09-1992	US 5256766 A	26-10-1993
		US 5688768 A	18-11-1997
		AU 665752 B	18-01-1996
		AU 1456892 A	15-09-1992
		CA 2104394 A	20-08-1992
		EP 0572553 A	08-12-1993
		JP 6508742 T	06-10-1994
		NZ 241666 A	26-07-1995
		US 6024936 A	15-02-2000
		US 5759994 A	02-06-1998
		US 5856448 A	05-01-1999
		US 5849507 A	15-12-1998
		US 6124101 A	26-09-2000
		US 5798248 A	25-08-1998
